

PSYCHOLOGICAL NOTES.

WILLIAM O. KROHN, PH. D.

Steps have been taken at Northwestern University to establish a laboratory in experimental psychology and an instructor or laboratory demonstrator will soon be appointed.

The work in pedagogy and psychology at the University of the State of Missouri, heretofore undertaken by one man has just been divided and an additional professor is to be appointed, who will devote his time entirely to psychology. This must of course ultimately lead to the founding of a well equipped psychological laboratory.

The change in administration in Illinois has made it necessary for Dr. Dewey to resign his position as superintendent of the Illinois Hospital for the Insane, at Kankakee—a position which he has so ably filled for fourteen years. Dr. Dewey has achieved his international reputation largely through his ardent advocacy of the “detached ward” or “cottage” system which he has brought to such a degree of perfection at the Kankakee hospital. As one result of his championship of this idea there are at this asylum excellent accommodations for 1000 male and 1000 female patients. Dr. S. V. Clevenger, so widely and so favorably known, is to be Dr. Dewey's successor. In such a change it is a delightful satisfaction to know that the position is to be held by a man who brings with him such a large amount of experience and skill as Dr. Clevenger certainly does. Dr. Dewey will locate in Chicago where he opens a private sanitarium for the treatment of nervous and mental diseases.

Hypnotism among the Insane.—By George M. Robertson, (Journal of Mental Science, January, 1893). Writer simply recounts result of observations made during visits at Nancy and Paris, the real work having been done by Voisin, Bernheim, and Luys. To give the article color of originality he refers to some experiments made at the Morningside Asylum which he uses Bernheim's method. R. acknowledges what others have known for years that the insane are especially difficult to hypnotize. Also with the insane the hypnotic sleep is of shorter duration than with the normal individual. Summarizing R's results it is found that hypnotism may be used among the insane:—

1. *As a therapeutic* in (a.) insomnia. Hypnotic sleep is more closely allied to healthy normal sleep than drugged sleep—therefore is of great service where brain nutrition is bad without depressing effect of drugs.

b. As a sedative in excitement.

c. To dispel temporary hallucinations.

2. *May be used for purposes of management.*

a. To overcome the morbid resistance of patients. Patients often refuse to do what is necessary for their highest good, and by hypnotizing them they can be made to do what is desired.

b. As a substitute for restraint.

Neuroses Convulsives et Affaiblissement Intellectuel.—By Prof. X. Francotte, (Bulletin de la Société de Médecine Mentale de Belgique, December, 1892). The association of convulsive phenomena with psychical disturbances is very common, but it is still a question as to whether there is any causal relation between them. From three interesting cases which have been carefully observed by Prof. F., the following summary may be made.

All three of these cases have a common character:—chronic convulsive neurosis accompanied by intellectual enfeeblement ending in insanity. Is it not possible, he says, to attribute the intellectual weakening to the convulsive condition? His theory is based on Ribot's "Psychology of attention," which demonstrates from numerous examples, that a great expenditure of motion and the condition of attention are antagonistic. Every one knows that reflection is impossible while running, or during any intense movement. On the other hand, when an audience is deeply attentive, it is said, according to the French, "*on y entendrait voler une mouche*," or as our localism gives it, "one could hear a pin drop." How quickly a flutter of movement passes over an audience after a period of strained attention.

Prof. F., concludes then, that chronic convulsive conditions, to say the least, favor and never inhibit intellectual degeneracy.

Visual Cerebral Centres in the case of the Dog and the Monkey.—By Vitzou, (2. Congr. de Physiol. tem. à Liège, an 1892). After having anæsthetized a dog by atropo-morphine and chloroform, the posterior part of the left hemisphere of the brain was extirpated. Awakening the animal, and placing a bandage over the eye corresponding to the injured side, the sight was indistinct, and he stumbled over obstacles. He took no notice of a